

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68112 A2

(51) International Patent Classification⁷: **A61K 38/00**

(21) International Application Number: PCT/EP01/02882

(22) International Filing Date: 14 March 2001 (14.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/189,091 14 March 2000 (14.03.2000) US

(71) Applicants and

(72) Inventors: **GÖKE, Burkhard** [DE/DE]; Römerstrasse
47 a, 82131 Gauting (DE). **SCHIRRA, Jörg** [DE/DE];
Meyerbeerstrasse 4, 81247 München (DE).

(74) Agent: **DÖRRIES, FRANK-MOLNIA & POHLMAN**;
Postfach 221661, 80538 Munich (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: EFFECTS OF GLUCAGON-LIKE PEPTIDE-1 (7-36) ON ANTRO-PYLORO-DUODENAL MOTILITY

(57) Abstract: The present invention provides an effective method for inhibiting antro-duodenal motility in healthy subjects and patients suffering from various disorders, without the side effects associated with other pharmaceutical compositions. GLP-1 slows antro-duodenal motility and may be used for the treatment or prevention of gastrointestinal disorders such as diarrhea, postoperative dumping syndrome and irritable bowel syndrome, and also premedication in endoscopic procedures.

WO 01/68112 A2

EFFECTS OF GLUCAGON-LIKE PEPTIDE-1 (7-36) ON ANTRO-PYLORO-DUODENAL MOTILITY

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to inhibiting antro- duodenal motility with GLP-1 and methods to alleviate discomfort during endoscopy and to alleviate symptoms of gastrointestinal disorders.

2. Description of the Related Art

Glucagon has been widely used to cause a variable reduction in gastroduodenal motility. The effect of glucagon appears to be dose-dependent with a minimally effective dose being 0.5 mg. Glucagon, however, does not facilitate colonoscopic evaluation (Norfleet, Gastrointest. Endosc., 24, 164-5, 1978), and at doses as high as 2 mg glucagon does not reduce contractions in the antrum (Gregerson et al., Scand. J. Gastroenterol. 23 (Supp 152), 42-47 (1988)). Furthermore, glucagon is contraindicated in persons with diabetes (Paul & Freyschmidt, ROFO Rortschr. Geb. Rontgenstr. Nuklearmed., 125, 31-7 (1996)), is expensive and its efficacy has been questioned.

Side effects associated with the use of glucagon include nausea and vomiting. The effects are dose-dependent and can appear at a dose of 1 mg (Larsen et al., Scand. J. Gastroenterol. 21, 634-640, 1986; Gregersen et al., supra, Diamant Handbook Experimental Pharm, Lefevre ed., Vol. 66/2, 611-643, 1983). As dosages required to sufficiently reduce motility frequently exceed 1 mg, side effects from glucagon use are common. Such side effects

render the patient extremely uncomfortable and often cause the endoscopic procedure to be interrupted or aborted.

Hyoscamine sulfate has antispasmodic activity and been used for treatment of irritable bowel syndrome and also has adverse side effects (Lahoti *et al.*, *Gastrointest. Endosc.* 46, 139-142 (1997)). Otreocide, a somatostatin analog, has also been used and proven effective in treating clinically significant diarrhea during rapid detoxification and postoperative dumping syndrome, but is associated with an unacceptable incidence of bradycardia; its long-term use is limited by side effects.

The proglucagon-derived glucagon-like peptide-1(7-36)amide (GLP-1) is a gastrointestinal hormone that is released postprandially from the L-cells of the gut (Göke *et al.*, *Eur. J. Clin. Invest.* 21, 135-44 (1991); Schirra *et al.*, *J. Clin. Invest.* 97, 92-103 (1996)). Previous studies in humans have shown synthetic GLP-I to substantially retard gastric emptying of liquid and solid meals (Schirra *et al.*, *J. Endocrinol.* 156, 177-86 (1998); Wettergren *et al.*, *Dig. Dis. Sci.* 38, 665-673 (1993); Schirra *et al.*, *Proc. Assoc. Am. Physicians* 109, 84-97 (1997)). Transpyloric pulsatile flow regulated by the motility of the antro-pyloro-duodenal region is a major mechanism of gastric emptying (Malbert & Mathis, *Gastroenterol.* 107, 37-46 (1994); Anvari *et al.*, *J. Physiol. (London)* 488, 193-202 (1995)). Antral contractions, and especially antro-duodenal coordinated ones, were shown to be associated with the gastric emptying rate of liquids (Schirra *et al.*, *J. Clin. Invest.* 97, 92-103 (1996); Camilleri *et al.*, *Am J Physiol* 249, G580-585 (1985); Houghton *et al.*, *Gastroenterol.* 94, 1276-84 (1988)) and solids (Fraser *et al.*, *Am. J. Physiol.* 264, G195-201 (1993)). Tonic and localized phasic pressure increases generated by the pylorus provide an important braking mechanism, diminishing gastric outflow (Anvari *et al.*, *J. Physiol. (London)* 488, 193-202 (1995); Heddl *et al.*, *Dig. Dis. Sci.* 38, 856-69 (1993); Heddl *et al.*, *Gut* 29, 1349-57 (1988); Tougas *et al.*, *Gut* 33, 466-471 (1992)).

SUMMARY

It is therefore an object of the present invention to provide a method for inhibiting antro-duodenal motility with an effective, therapeutic composition that has minimal side effects. It is well known that GLP-1 does not cause hypoglycemia – and it did not cause hypoglycemia during the experiments discussed in the present application, nor did it cause any other side effects. Accordingly, a dosage unit comprising a GLP-1 molecule and a pharmaceutically suitable excipient is also disclosed.

In a related vein, the present invention also encompasses a method for premedicating in endoscopic procedures, comprising administering a GLP-1 molecule prior to or during an endoscopic procedure.

Another embodiment of the present invention is a method for treating or preventing gastrointestinal disorders, including but not limited to, irritable bowel syndrome, non-infectious acute and chronic diarrhea and post-operative dumping syndrome, that comprises administering to the patient therapeutically effective amount of a GLP-1 molecule.

Further encompassed by this invention is a method for treating or preventing symptoms associated with narcotics withdrawal, by administering a GLP-1 molecule as described above.

In another embodiment, the invention includes a method for inhibiting pyloric motility in a patient in need thereof that comprises administering to the patient a therapeutically effective amount of an antagonist to a GLP-1 molecule.

DESCRIPTION OF FIGURES

Figure 1. Plasma immunoreactivities of GLP-1 in response to intravenous infusions of saline, GLP-1(7-36)amide at 0.4 and 1.2 pmol·kg⁻¹·min⁻¹ with concomitant duodenal perfusion of saline or lipid (2.5 kcal/min) in eleven healthy volunteers. Mean±SEM. For statistical analysis, see Table 3.

Figure 2. Contraction frequencies (upper panel) and amplitudes (lower panel) in the antrum and duodenum in response to intravenous infusions of saline, GLP-1(7-36)amide at 0.4 and 1.2 pmol·kg⁻¹·min⁻¹ during concomitant duodenal perfusion of saline (A) or lipid (2.5 kcal/min, B) in eleven healthy volunteers. Mean±SEM. For statistical analysis, see Table 1.

Figure 3. Isolated pyloric pressure waves (A) and pyloric tone (B) in response to intravenous infusions of saline, GLP-1(7-36)amide at 0.4 and 1.2 pmol·kg⁻¹·min⁻¹ during concomitant duodenal perfusion of saline (upper panel) or lipid (2.5 kcal/min, lower panel) in eleven healthy volunteers. Mean ± SEM. *:p<0.05 for comparison of time points indicated by arrows (paired t-test). For further statistical analysis, see Table 2.

Figure 4. Occurrence of duodenal phase III-like activity with duodenal perfusion of saline (upper panel) or lipid (2.5 kcal/min, lower panel) in eleven healthy volunteers. In seven of eleven subjects an activity front was seen within 10 min. after start of low dose GLP-1 in the interdigestive state or duodenal lipid perfusion in the postprandial studies, respectively. Length of solid bars represents length of contraction burst.

Figure 5. Manometric tracings showing the effects of intravenous infusion of GLP-1 at 0.4 pmol·kg⁻¹·min⁻¹ during duodenal perfusion of saline (A) and lipid (2.5 kcal/min, B). In the interdigestive state (A), GLP-1 immediately inhibits antroduodenal motility, and induces a sustained increase of basal pyloric pressure with a concomitant short lasting stimulation of IPPWs. During duodenal lipid perfusion (B), antral and duodenal contractility are abolished with GLP-1, and basal pyloric pressure further increases in addition to the effect of lipid alone paralleled by a stimulation of IPPWs.

Figure 6. Effects of intravenous infusions of saline, GLP-1(7-36)amide at 0.4 and 1.2 pmol·kg⁻¹·min⁻¹ on plasma glucose (A) and immunoreactivities of glucagon (B), insulin (C), and pancreatic polypeptide (D) during concomitant duodenal perfusion of saline or lipid (2.5 kcal/min) in eleven healthy volunteers. Mean±SEM of incremental values over basal. For statistical analysis, see Table 3.

DETAILED DESCRIPTION

Previous studies in humans have shown synthetic GLP-1 to substantially retard gastric emptying of liquid and solid meals (Schirra *et al.*, *J. Endocrinol.* 156, 177-86 (1998); Wettergren *et al.*, *Dig. Dis. Sci.* 38, 665-673 (1993); Schirra *et al.*, *Proc Assoc Am Physicians* 109, 84-97 (1997)). In addition to its insulinotropic and glucagonostatic action, reduction of the gastric emptying rate may be considerably involved in the glucose-lowering effect of GLP-1 in healthy subjects and patients with diabetes mellitus (Schirra *et al.*, *J. Endocrinol.* 156, 177-86 (1998); Schirra *et al.*, *Proc Assoc Am Physicians* 109, 84-97 (1997)). Independent of the tonic pressure generated by the proximal stomach, transpyloric pulsatile flow regulated by the motility of the antro-pyloro-duodenal region is a major mechanism of gastric emptying (Malbert & Mathis, *Gastroenterol.* 107, 37-46 (1994); Anvari *et al.*, *J. Physiol. (London)* 488, 193-202 (1995)). Antral contractions, and especially antro-duodenal coordinated ones, were shown to be associated with the gastric emptying rate of liquids (Schirra *et al.*, *J. Clin. Invest.* 97, 92-103 (1996); Camilleri *et al.*, *Am J Physiol* 249, G580-585 (1985); Houghton *et al.*, *Gastroenterol.* 94, 1276-84 (1988)) and solids (Fraser *et al.*, *Am. J. Physiol.* 264, G195-201 (1993)). Tonic and localized phasic pressure increases generated by the pylorus provide an important braking mechanism diminishing gastric outflow (Anvari *et al.*, *J. Physiol. (London)* 488, 193-202 (1995); Heddle *et al.*, *Dig. Dis. Sci.* 38, 856-69 (1993); Heddle *et al.*, *Gut* 29, 1349-57 (1988); Tougas *et al.*, *Gut* 33, 466-471 (1992)).

The present application describes the effects of graded doses of synthetic GLP-1 on the motility of the antro-pyloro-duodenal region during the interdigestive and postprandial state in human, the latter being elicited by duodenal perfusion of lipid. Duodenal perfusion of lipid was used, instead of oral meal ingestion, to provide a constant duodenal nutrient load independent of gastric emptying. This particular meal was chosen to establish a stable postprandial motility pattern, and in order to minimize plasma glucose and insulin excursions. Finally, the effects of GLP-1 were compared with those of lipid, the classical physiological stimulator of pyloric motility.

The applicants have discovered that GLP-1 provides a cost-effective therapeutic composition for use in preventing or treating various gastrointestinal disorders. Furthermore,

GLP-1 can be used to premedicate for endoscopy and treat symptoms of narcotics withdrawal. Other medications do not share the same effectiveness as GLP-1 and are often accompanied by adverse side effects. Such side effects include, but are not limited to, nausea and vomiting. These side effects have not been observed in LP-1 treated patients. [

5 Assessing gastric motility

Inhibition of antro-duodenal activity is assessed directly by motility recordings using methods that are well known to the skilled artisan. Contractile events are analyzed with a computer and validated software (Katschinski *et al.*, *Gastroenterol.* 103, 383-91 (1992)) and only peaks with amplitudes of at least 10 mmHg and a duration of at least 2 s are considered true
10 contractions. Duodenal phase III is defined as the occurrence of regular contractions at a frequency of $\geq 10/\text{min}$ for at least 2 min in the duodenum propagated aborally.

In one embodiment, the data are analyzed in 10-min segments separately for antrum and duodenum by summarizing (frequency, motility index) or averaging (amplitude) the values derived from, the two antral and three duodenal side holes, respectively. Motility indexes are
15 identified as the area under the contractions and are expressed in $\text{mmHg}\cdot\text{s}\cdot\text{min}^{-1}$. Antral contractions are determined to be antro-pyloro-duodenally propagated waves if the onset of the pressure wave recorded in the most proximal duodenal side hole, occurs within 5 sec after the onset of a pressure wave recorded in one of the antral side holes and if both wave are registered by all side holes in between.

As used herein, "inhibition" of antro-pyloro-duodenal motility is defined as a reduction in the motility index. Other methods for measuring motility can be used. One of skill in the art will appreciate that a significant inhibition, including total inhibition, of motility will be useful. Pyloric tone can be calculated as change from basal, the latter being determined as mean pyloric tone during the basal period before starting the experiments.
20

25

GLP-1 Molecules

As used herein, a "GLP-1 molecule" includes the following. Mammalian GLP peptides and glucagon are encoded by the same gene. In the ileum the phenotype is processed into two major classes of GLP peptide hormones, namely GLP-1 and GLP-2. GLP-1(1-37) has the
30 sequence His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO:1). GLP-1

(1-37) is amidated post-translationally to yield GLP-1(1-36)NH₂, which has the sequence His Asp Glu Phe Glu Arg His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH₂) (SEQ ID NO:2); or is enzymatically processed to yield GLP-1(7-37), which has the sequence His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO:3). GLP-1(7-37) can also be amidated to yield GLP-1(7-36)amide, which is the natural form of the GLP-1 molecule, and which has the sequence His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH₂) (SEQ ID NO:4). Likewise, GLP-1(1-36)(NH₂) can be processed to GLP-1(7-36)(NH₂).

Intestinal L cells secrete GLP-1(7-37) (SEQ ID NO:3) and GLP-1(7-36)NH₂ (SEQ ID NO: 4) in a ratio of 1 to 5, respectively. These truncated forms of GLP-1 have short half-lives *in situ*, i.e., less than 10 minutes, and are inactivated by an aminodipeptidase IV to yield Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO:5); and Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH₂) (SEQ ID NO:6); respectively. The peptides Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly (SEQ ID NO:5) and Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg (NH₂) (SEQ ID NO:6), have been speculated to affect hepatic glucose production, but do not stimulate production or release of insulin from the pancreas.

As used in this specification, the term "GLP-1 molecule" includes GLP-1(1-37), GLP-1(1-36)NH₂, GLP-1(7-37), GLP-1(7-36)NH₂ ("GLP-1(7-36)amide") (collectively referred to as "GLP-1 peptides"). The present invention includes the use of recombinant human GLP-1 peptides as well as GLP-1 peptides derived from other species, whether recombinant or synthetic.

"GLP-1 molecule" further denotes biologically active variants, analogs, and derivatives of GLP-1 peptides. "Biologically active," in this context, means having GLP-1(7-36) biological activity, but it is understood that the activity of the variant can be either less potent or more potent than native GLP-1(7-36)amide. GLP-1(7-36)amide is a native, biologically active form of GLP-1. See Göke & Byrne, *Diabetic Medicine*, 13, 854-860 (1996). GLP-1 molecules of the

present invention include polynucleotides that express agonists of GLP-1, i.e. activators of the GLP-1 receptor molecule and its secondary messenger activity found on, *inter alia*, insulin-producing β -cells. GLP-1 mimetics that also are agonists of β -cells include, for example, chemical compounds specifically designed to activate the GLP-1 receptor.

5 GLP-1 molecule biological activity can be determined by *in vitro* and *in vivo* animal models and human studies as is well known to the skilled artisan. Included as GLP-1 molecules are any molecules, whether they be peptides, peptide mimetics, or other molecules that bind to or activate a GLP-1 receptor, such as the GLP-1(7-36)amide receptor, and its second messenger cascade. GLP-1 receptors are cell-surface proteins found, for example, on insulin-producing
10 pancreatic β -cells. The GLP-1(7-36) receptor has been characterised in the art. Methods of determining whether a chemical or peptide binds to or activates a GLP-1 receptor are known to the skilled artisan and are preferably carried out with the aid of combinatorial chemical libraries and high throughput screening techniques. GLP-1 molecules include species having insulinotropic activity and that are agonists of, i.e. activate, the GLP-1 receptor molecule and its
15 second messenger activity on, *inter alia*, insulin-producing β -cells.

 GLP-1 biological activity can be determined by standard methods, in general, by receptor-binding activity screening procedures which involve providing appropriate cells that express the GLP-1 receptor on their surface, for example, insulinoma cell lines such as RINmSF cells or INS-1 cells. See also Mojsov, Int J Pept Protein Res 40, 333-43 (1992)
20 and EPO708170A2. Cells that are engineered to express a GLP-1 receptor also can be used. In addition to measuring specific binding of tracer to membrane using radioimmunoassay methods, cAMP activity or glucose dependent insulin production can also be measured. In one method, a polynucleotide encoding the receptor of the present invention is employed to transfect cells to thereby express the GLP-1 receptor protein. Thus, for example, these methods may be employed
25 for screening for a receptor agonist by contacting such cells with compounds to be screened and determining whether such compounds generate a signal, i.e activate the receptor.

 Polyclonal and monoclonal antibodies can be utilized to detect purify and identify GLP-1 like peptides for use in the methods described herein. Antibodies such as ABGA1178 detect intact unprocessed GLP-1(1-37) or N-terminally-truncated GLP-1(7-37) or (7-36)amide. Other
30 antibodies detect on the very end of the C-terminus of the precursor molecule, a procedure which allows by subtraction to calculate the amount of biologically active truncated peptide, i.e. GLP-

1(7-37)amide (Orskov et al., *Diabetes*, 42, 658-661 (1993); Orskov et al., *J. Clin. Invest.* 87, 415-423 (1991)).

Other screening techniques include the use of cells which express the GLP-1 receptor, for example, transfected CHO cells, in a system which measures extracellular pH or ionic changes caused by receptor activation. For example, potential agonists may be contacted with a cell which expresses the GLP-1 protein receptor and a second messenger response, e.g. signal transduction or ionic or pH changes, may be measured to determine whether the potential agonist is effective.

Agonists of glucagon-like peptide that exhibit activity through the GLP-1(7-36)amide receptor have been described. See EP 0708179 A2; Hjorth et al., *J. Biol. Chem.* 269; 30121 (1994); Siegel et al., Amer. Diabetes Assoc. 57th Scientific Session, Boston (1997); Hareter et al., Amer. Diabetes Assoc. 57th Scientific Session, Boston (1997); Adelhorst et al., *J. Biol. Chem.* 269, 6275 (1994); Deacon et al., 16th International Diabetes Federation Congress Abstracts, *Diabetologia Supplement* (1997); Irwin et al., *Proc. Natl. Acad. Sci. USA* 94; 7915 (1997); Mojsov, *Int. J. Peptide Protein Res.* 40; 333 (1992). Göke & Byrne, *Diabetic Medicine* 13; 854 (1996). Recent publications disclose Black Widow GLP-1 and Ser² GLP-1. See Holz & Hakner, *Comp. Biochem. Physiol., Part B* 121; 177 (1998) and Ritzel et al., *J. Endocrinol* 159; 93 (1998).

"GLP-1 molecules" also include peptides that are encoded by polynucleotides that express biologically active GLP-1 variants as defined herein. Also included in the present invention are GLP-1 molecules that are peptides containing one or more amino acid substitutions, additions or deletions, compared with GLP-1 (7-36) amide. In one embodiment, the number of substitutions, deletions, or additions is 30 amino acids or less, 25 amino acids or less, 20 amino acids or less, 15 amino acids or less, 10 amino acids or less, 5 amino acids or less or any integer in between these amounts. In one aspect of the invention, the substitutions include one or more conservative substitutions. A "conservative" substitution denotes the replacement of an amino acid residue by another, biologically active similar residue. Examples of conservative substitution include the substitution of one hydrophobic residue, such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acids, or glutamine for

asparagine, and the like. The following table lists illustrative, but non-limiting, conservative amino acid substitutions.

<u>ORIGINAL RESIDUE</u>	<u>EXEMPLARY SUBSTITUTIONS</u>
ALA	SER, THR
ARG	LYS
ASN	HIS, SER
ASP	GLU, ASN
CYS	SER
GLN	ASN, HIS
GLU	ASP, GLU
GLY	ALA, SER
HIS	ASN, GLN
ILE	LEU, VAL, THR
LEU	ILE, VAL
LYS	ARG, GLN, GLU, THR
MET	LEU, ILE, VAL
PHE	LEU, TYR
SER	THR, ALA, ASN
THR	SER, ALA
TRP	ARG, SER
TYR	PHE
VAL	ILE, LEU, ALA
PRO	ALA

- 5 It is further understood that GLP-1 peptide variants include the above described peptides, which have been chemically derivatized or altered, for example, peptides with non-natural amino acid residues (e.g., taurine residue, β - and γ -amino acid residues and D-amino acid residues), C-terminal functional group modifications, such as amides, esters, and C-terminal ketone

modifications and N-terminal functional group modifications, such as acylated amines, Schiff bases, or cyclization, such as found, for example, in the amino acid pyroglutamic acid.

Also included in the present invention are peptide sequences having greater than 50 percent sequence identity, and preferably greater than 90 percent sequence identity to (1) SEQ ID NOS:1, 2, 3, 4; and (2) to truncated sequences thereof. As used herein, sequence identity refers to a comparison made between two molecules using standard algorithms well known in the art. The preferred algorithm for calculating sequence identity for the present invention is the Smith-Waterman algorithm, where SEQ ID NO:1 is used as the reference sequence to define the percentage identity of polynucleotide homologs over its length. The choice of parameter values for matches, mismatches, and inserts or deletions is arbitrary, although some parameter values have been found to yield more biologically realistic results than others. One preferred set of parameter values for the Smith-Waterman algorithm is set forth in the "maximum similarity segments" approach, which uses values of 1 for a matched residue and -1/3 for a mismatched residue (a residue being either a single nucleotide or single amino acid) (Waterman, Bull.Math. Biol. 46, 473-500 (1984)). Insertions and deletions (indels), x, are weighted as

$$x_k = 1 + k/3,$$

where k is the number of residues in a given insert or deletion (Id.).

For instance, a sequence that is identical to the 42 amino acid residue sequence of SEQ ID NO:1, except for 18 amino acid substitutions and an insertion of 3 amino acids, would have a percent identity given by:

$$\begin{aligned} & [(1 \times 42 \text{ matches}) - (\alpha \times 18 \text{ mismatches}) \\ & - (1 + 3/3 \text{ indels})] / 42 = 81\% \text{ identity} \end{aligned}$$

Also included in "GLP-1 molecules" of the present invention are six peptides in Gila monster venoms that are homologous to GLP-1. Their sequences are compared to the sequence of GLP-1 in Table 1.

TABLE I

Position 1
a. HAEGTFTSDVSSYLEGQAAKEFIAWLVKGR(NH₂)

- b. HSDGTFTSDLSKQMEEEEAVRLFIEWLKNNGPSSGAPPPS (NH₂)
 c. DLSKQMEEEEAVRLFIEWLKNNGPSSGAPPPS (NH₂)
 d. HGEGTFTSDLSKQMEEEEAVRLFIEWLKNNGPSSGAPPPS (NH₂)
 e. HSDATFTA EYSKLLAKLALQKYLE SILGSSTSPRPPSS
 5 f. HSDATFTA EYSKLLAKLALQKYLE SILGSSTSPRPPS
 g. HSDAIFTEEYSKLLAKLALQKYLASILGSRTSPPP (NH₂)
 h. HSDAIFTQ QYSKLLAKLALQKYLASILGSRTSPPP (NH₂)

a = GLP-1(7-36)amide (SEQ. ID NO: 4)

10 b = exendin 3 (SEQ. ID NO: 7).

c = exendin 4 (9-39(NH₂)) (SEQ. ID NO: 8).

d = exendin 4 (SEQ. ID NO: 9).

e = helospectin I (SEQ. ID NO: 10).

f = helospectin II (SEQ. ID NO: 11).

15 g = helodermin (SEQ. ID NO: 12).

h = Q⁸, Q⁹ helodermin (SEQ. ID No:13).

Peptides (a, b, d, e, f and g) are homologous in positions 1, 7, 11 and 18. GLP-1 and
 20 exendins are further homologous in positions, 4, 5, 6, 8, 9, 15, 22, 23, 25, 26 and 29. In position
 2, A, S and G are structurally similar. In position 3, residues D and E (Asp and Glu) are
 structurally similar. In positions 22 and 23, F (Phe) and I (Ile) are structurally similar to Y (Tyr)
 and L (Leu), respectively. Likewise, in position 26, L and I are structurally equivalent. Thus, of
 the 30 residues of GLP-1, exendins 3 and 4 are identical in 15 positions and equivalent in 5
 25 additional positions. The only positions where radical structural changes are evident are at
 residues 16, 17, 19, 21, 24, 27, 28 and 30. Exendins also have 9 extra residues at the carboxyl
 terminus.

The GLP-1 molecules of the invention that are peptides that can be made by solid state
 chemical peptide synthesis. Such peptides can also be made by conventional recombinant
 30 techniques using standard procedures described in, for example, Sambrook & Maniatis.
 "Recombinant", as used herein, means that a gene is derived from a recombinant (e.g., microbial
 or mammalian) expression system which has been genetically modified to contain
 polynucleotide encoding a GLP-1 molecule as described herein.

The GLP-1 like peptides can be recovered and purified from recombinant cell cultures by methods including, but not limited to, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance liquid chromatography (HPLC) can be employed for final purification steps.

The GLP-1 molecule peptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from prokaryotic or eukaryotic hosts (for example by bacteria, yeast, higher plant, insect and mammalian cells in culture or *in vivo*). Depending on the host employed in a recombinant production procedure, the polypeptides of the present invention are generally non-glycosylated, but may be glycosylated.

Particularly preferred GLP-1 molecules of the invention are GLP-1(7-36)amide, and GLP-1(7-37) and exendin-4.

Synthetic GLP-1(7-36)amide can be purchased and delivered with a peptide content of 87.1% and a peptide purity of >99%. According to known methods in the field, the peptide can be dissolved in 1% human serum albumin, filtered through 0.2 µm nitrocellulose filters and then stored at -70°C [13]. Samples are tested for pyrogens and bacterial growth standard techniques.

Without being bound by a particular mechanism, the present inventor suggests that the following motor effects of GLP-1 are important mediators of reduced antro-duodenal motility: (i) inhibition of antral waves in general, (ii) reduction of transpyloric propagated antral waves in particular and (iii) simultaneous stimulation of localized phasic and tonic pyloric contractions. In dogs, a similar motor pattern has been demonstrated to be associated with reduction of transpyloric flow and inhibition of gastric emptying of a non-caloric liquid meal with infusion of synthetic GLP-1 (Anvari *et al.*, *Dig. Dis. Sci.* 43, 1133-40 (1998)). In human, subcutaneously injected GLP-1 dose-dependently inhibits antral and coordinated antroduodenal contractions parallel to prolongation of the gastric emptying lag phase of a mixed liquid meal (Schirra *et al.* *Proc. Assoc. Am. Physicians* 109, 84-97 (1997)). Additionally, preliminary data obtained from human studies indicate a strong relaxation of the proximal stomach in response to intravenous GLP-1 using the same dosages as in the present study (Wank *et al.*, *Gastroenterol.* 114, A1190

(abstract) (1998)). Thus, exogenous GLP-1 reduces driving forces and stimulates braking mechanisms of gastric outflow, thereby impacting on all motor sites known to determine gastric emptying.

5

Formulating a dosage unit

Another embodiment of the present invention is a dosage unit for use in inhibiting antro-duodenal motility in a patient comprising GLP-1 and a pharmaceutically suitable excipient is described. Preferably, the dosage unit is in the range of about 0.4 to 2.4 pmol·kg⁻¹·min⁻¹. Still
10 more preferably, the dosage unit is in the range of about 0.8 to 1.2 pmol·kg⁻¹·min⁻¹. For the purpose of this invention, the term "about" is defined as +/- 10%. For example, a dosage unit in the range of about 0.4 to 2.4 pmol·kg⁻¹·min⁻¹ means 0.196 to 2.64 pmol·kg⁻¹·min⁻¹.

The composition of the present invention can be used as a systematic or local application by oral or parenteral administration. Alternatively, the composition may be applied as an
15 intravenous or subcutaneous injection. For use by the physician or patient, the composition may be provided in a dosage unit form containing an amount of a GLP-1 molecule, for example, a GLP-1(7-36)amide, with or without another antimotility agent. This will be effective in one or multiple doses for use in the inhibition of antro-duodenal motility. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including
20 the age and weight of the patient, the patient's physical condition and other factors. Preferably, a GLP-1 molecule is administered intravenously and the dosage unit is in the range of about 0.4 to 2.4 pmol·kg⁻¹·min⁻¹. Still more preferably, the dosage unit is in the range of about 0.8 to 1.2 pmol·kg⁻¹·min⁻¹.

The solid composition for oral administration of the present invention includes tablets,
25 preparations, granules and the like. In such a solid composition, one or more active ingredients may be mixed with at least one inactive diluent, for example, lactose, mannitol, glucose, hydroxypropyl cellulose, microcrystalline cellulose, starch, polyvinyl pyrrolidone, magnesium aluminate metasilicate and the like. According to the usual work-up, the composition may contain additives other than inactive diluent, for example, lubricant such as magnesium stearate;
30 disintegrant such as fibrous calcium gluconate; stabilizer such as cyclodextrin, for example, α,β- or γ-cyclodextrin; etherified cyclodextrin such as dimethyl-α-, dimethyl-β-, trimethyl-β-, or

hydroxypropyl- β -cyclodextrin; branched cyclodextrin such as glucosyl-, maltosyl-cyclodextrin; formylated cyclodextrin, cyclodextrin containing sulfur; phospholipid and the like. When the above cyclodextrins are used, inclusion compound with cyclodextrins may be sometimes formed to enhance stability. Alternatively, phospholipid may be sometimes used to form liposome, resulting in enhanced stability.

Tablets or pills may be coated with film soluble in the stomach or intestine such as sugar, gelatin, hydroxypropyl cellulose, hydroxypropylmethyl cellulose phthalate as needed. Further, they may be formed as capsules with absorbable substances such as gelatin.

A liquid composition for oral administration may contain pharmaceutically acceptable emulsion, solution, suspension, syrup, elixer as well as generally used inactive diluent. Such composition may contain, in addition to the inactive diluent, adjuvants such as lubricants and suspensions, sweetening agents, flavoring agents, preservatives, solubilizers, anti-oxidants and the like. The details of the additives may be selected from those described in any general textbook in the pharmaceutical field. Such liquid compositions may be directly enclosed in soft capsules.

Solutions for parenteral administration, for example, suppository, enema and the like according to the present invention include sterile, aqueous or non-aqueous solution, suspension, emulsion, detergent and the like. The aqueous solution and suspension includes, for example, distilled water, physiological saline and Ringer's solution.

The non-aqueous solution and suspension include, for example, propylene glycol, polyethylene glycol, fatty acid triglyceride, vegetable oil such as olive oil, alcohols such as ethanol, polysorbate and the like. Such composition may contain adjuvants such as preservatives, wetting agent, emulsifier, dispersant, anti-oxidants and the like.

Inhibiting antro- duodenal motility

The present invention encompasses a method for inhibiting antro-duodenal motility in a patient in need thereof, which comprises administering to the patient a composition comprising a GLP-1 molecule in a therapeutically effective amount. In a preferred embodiment, said GLP-1 molecule is GLP-1(7-36)amide. As described throughout the specification, the GLP-1 molecule can be administered using a variety of methods. Furthermore, as described herein, there are a

variety of patients that will benefit from the inhibition of antro-duodenal motility, using the inventive methods. The term "patient" as used herein refers to any mammal, including a human.

Premedicating with GLP-1

Included in this invention is a method for premedicating in endoscopic procedures, which
5 comprises administering to the patient a composition comprising a GLP-1 molecule in a therapeutically effective amount. Premedication is intended to reduce a patient's discomfort in endoscopic procedures and the like, by reducing antro-duodenal motility. Contraction or spasm of gastrointestinal smooth muscle imposes a technical obstacle, which must frequently be overcome in order to enable the clinician to successfully, perform endoscopic procedures.
10 "Premedication" includes administration of a GLP-1 molecule just prior to, or during the procedure. "Just prior to" means up to about 1 hour before beginning the procedure or before scope insertion, and "during" means any time during the procedure, including at cecal intubation and before or during scope withdrawal. Just prior to includes administration of a GLP-1 molecule up to about 45, 30, 20, 15, 10, 5 minutes, or 1 minute before beginning the procedure
15 or before scope insertion. . The term "endoscope" or "scope" is used to refer to a colonoscope, gastroscope, enteroscope, cytoscope or other types of medical endoscopes.

The term "endoscopic procedures" refers to those diagnostic procedures that utilize an instrument that is introduced into the gastrointestinal tract to provide direct visualization of the gastrointestinal tract, for examination and therapeutic purposes. Such purposes include direct
20 visualization, biopsy, access to the common bile duct, fluid aspiration and removal of foreign bodies, polyps, and other lesions. An example of a particular endoscopic procedure is esophagogastro-duodenoscopy, which is utilized for examination of the esophageal lumen, stomach and duodenum. Another example, endoscopicretrograde cholangiopancreatography (ERCP), enables visualization of the pancreatic duct, common bile duct and the entire biliary
25 tract, including the gall bladder. Further examples of endoscopic procedures are colonoscopy, sigmoidoscopy and barium enema examinations.

Treating and preventing gastrointestinal disorders

Another facet of this invention is a method for treating or preventing non-infectious acute
30 and chronic diarrhea in a patient in need thereof which comprises administering to the patient a composition comprising a GLP-1 molecule in a therapeutically effective amount. "Acute" is

defined as a short and relatively severe course and "chronic" means persisting over longer periods of time (Dorland's Medical Dictionary, 27th Edition). "Non-infectious" implies not caused by, or capable of being communicated by infection, as an infectious disease (Dorland's Medical Dictionary, 27th Edition).

- 5 In a related vein, the invention includes a method for treating or preventing postoperative "dumping syndrome" in a patient in need thereof, which comprises administering to the mammal a GLP-1 molecule in a therapeutically effective amount. Dumping syndrome is one of the most common causes of morbidity after gastric surgery and can occur in patients after damage to the vagal nerve in esophageal surgery (Vecht *et al*). It is characterized by both gastrointestinal and
- 10 vasomotor symptoms. Gastrointestinal symptoms include postprandial fullness, crampy abdominal pain, nausea, vomiting and explosive diarrhea. Vasomotor symptoms include diaphoresis, weakness, dizziness, flushing, palpitations, and an intense desire to lie down. Patients with severe dumping symptoms may limit their food intake to minimize symptoms and as a result lose weight and become malnourished. In severe cases, as a last resort surgical
- 15 treatment of dumping syndrome has been utilized.

- Pharmaceutical treatment for severe dumping includes octreotide acetate (Sandoz), a long acting somatostatin analog, which has been used with some success. Octreotide is administered subcutaneously and acts to slow gastric emptying, inhibit insulin release, and decrease enteric peptide secretion. Octreotide, however, is accompanied by several complications, including
- 20 injection site pain, tachyphylaxis, iatrogenic diabetes, malabsorption and cholelithiasis.

In another embodiment, the invention includes a method for treating or preventing irritable bowel syndrome (IBS) in a patient in need thereof, which comprises administering to the patient a GLP-1 molecule in a therapeutically effective amount.

- IBS is believed to be a heterogeneous group of disorders, characterized by chronic lower
- 25 gastrointestinal symptoms not associated with an identifiable organic cause. The "spastic" variety of IBS, which is characterized by abdominal pain and constipation, is believed to be more common than IBS associated with chronic diarrhea and postprandial urgency. However, as noted above, the "diarrhea predominance" variety is also common.

- The definition and diagnosis of IBS have been somewhat controversial, but attempts have
- 30 been made in recent years to reach some consensus regarding the definition of IBS and other functional or non-organic GI disorders, and the clinical criteria for their diagnosis (Drossman *et*

al., supra; Hasler et al., supra; Camilleri et al., supra; Drossman et al., supra; Thompson et al., 1989, Gastroenterol. Intl. 2:92-95; Manning et al., 1978, Br. Med. J. 2:653-654 and Thompson et al., supra). According to the so-called "Rome criteria," patients afflicted with chronic diarrhea not attributable to an organic cause and not accompanied by significant abdominal pain are
5 classified as having functional diarrhea, and those patients afflicted with chronic diarrhea not attributable to an organic cause and accompanied by significant abdominal discomfort are classified as having IBS with diarrhea predominance (Drossman et al., supra; Hasler et al., supra; Camilleri et al., supra; Drossman et al., supra; Thompson et al., supra; Manning et al., supra).

In another aspect, the invention is directed to a method for treating or preventing
10 symptoms associated with narcotics withdrawal in a patient in need thereof, which comprises administering to the patient a GLP-1 molecule in a therapeutically effective amount. Clinically significant diarrhea is commonly seen during the acute phase of withdrawal in persons addicted to narcotics such as heroin and methadone. Loose stool and diarrhea frequently accompany the acute reaction of withdrawal from opioid drugs. The problem becomes even more severe when
15 patients are sedated during the procedure, as they lose bowel control.

Assessing pyloric motility

The present invention also relates to therapeutic methods associated with stimulating pyloric motility. Two parameters used to describe pyloric motility are: 1) Isolated pyloric
20 pressure waves (IPPW), defined as contractions registered by the sleeve with or without simultaneous contractions in maximally one sleeve-side-hole in the absence of associated (± 3 sec) waves of any amplitude in the adjacent antral and duodenal side-holes and 2) pyloric tone. Pyloric tone is measured by deducting the basal pressure recorded by the antral side hole at the proximal end of the sleeve, from the basal pressure recorded by the sleeve. Basal pressures is
25 defined as the mean pressure after excluding contractions; it is obtained each minute and used to calculate mean pyloric tone.

The invention also includes a method for inhibiting pyloric motility in a patient in need thereof, which comprises administering to the patient a GLP-1 antagonist in a therapeutically effective amount. Also described in the examples, GLP-1 stimulated pyloric motility. The
30 effects of GLP-1 in relation to those of lipid, the classical physiological stimulator of pyloric motility, were compared.

A comparison of pyloric stimulation by GLP-1, with and without duodenal lipid, provides new insights in the regulation of pyloric motility. Pyloric tone stably increases with both duodenal lipid and exogenous GLP-1 and the tonic response to duodenal lipid can be further enhanced by exogenous GLP-1. GLP-1 can stimulate pyloric tone at least as strong as duodenal lipid, and the interaction of both was additive. Conversely, GLP-1 was not able to stimulate IPPWs as strong as duodenal lipid, even at supraphysiological levels. Thus, the phasic pyloric response to duodenal lipid may be less dependant on GLP-1 release, and the mechanisms for GLP-1 stimulation of pyloric motility may be different than that of lipid.

EXAMPLES

The following examples illustrate the present invention but are not in any way intended to limit the scope of this invention. The following abbreviations are used in the examples below.

IPPW	isolated pyloric pressure wave
PP	pancreatic polypeptide
CCK	cholecystokinin

Measurement of plasma hormone levels and GLP-1.

Plasma glucose concentrations were measured using a glucose analyzer by the glucose oxidase method (YSI 1500 G; Schlag, Bergisch-Gladbach, Germany), with a coefficient of variation of <2%. Plasma insulin was measured by the Abbott Imx Microparticle Enzyme Immunoassay, with an average intraassay coefficient of variation of 5%. Plasma immunoreactivities of C-peptide, glucagon, and pancreatic polypeptide (PP) were analyzed by commercially available radioimmunoassay kits (Biermann, Bad Nauheim, Germany and Euradiagnostica, The Netherlands [PP]). Immunoreactive (IR) GLP-1 was measured using the specific polyclonal antibody GA 1178 (Affinity Research, Nottingham, United Kingdom) as described previously [2]. The detection limit of the assay was 0.25 pmol/l. The antiserum did not crossreact with glucose-dependent insulintropic peptide (GIP), pancreatic glucagon, glicentin, oxyntomodulin, or GLP-2. Intra- and inter-assay coefficients of variation were 3.4 and 10.4%, respectively. All values were expressed as mean \pm SEM. Plasma and motility parameters

were separately analyzed for each 60 min period. Pyloric tone was calculated as change from basal, the latter being determined as mean pyloric tone during the total basal period before starting the experiments. Differences of plasma hormones and glucose compared to the basal state were calculated as integrated values over basal (area under the response curve; AUC).

5 Basal levels of plasma parameters were determined as the mean of the two basal values just before the start of each experiment. All samples were first tested for normality by the Kolmogoroff-Smirnoff test. Differences between experimental sets for each parameter were analysed by two-way repeated measures analysis of variance employing intravenous infusion and duodenal perfusion as factors. When this analysis indicated differences, a Student-Newman-Keuls multicomparison test was performed. Differences were considered significant at $P < 0.05$.

10

Example 1. *Motility recordings and blood samples were taken from human patients*

Eleven healthy male volunteers, 23 to 28 years old and within 10% of ideal body weight, participated in the studies. None of them were taking medication, or suffering from gastrointestinal symptoms or systemic disease.

15

All studies were performed after an overnight fast. Two experiments were performed, separated by at least 1 week on subjects lying in a semirecumbent position. An indwelling catheter was inserted into an antecubital vein for intravenous infusions. A second catheter was inserted into a dorsal vein of the contralateral hand in a retrograde fashion for sampling of arterialized blood (Schirra *et al. Proc Assoc Am Physicians* 109, 84-97 (1997)).

20

Each experiment started after a basal period of at least 30 min, of which the last 15 min displayed a low motor activity (< 5 antral contractions/10 min). On the interdigestive study day, 0.154 M saline was continuously perfused into the region of the lower duodenal flexure at a rate of 2.5 ml/min. In order to obtain physiological postprandial and supraphysiological plasma levels, a 60 min period of intravenous saline infusion was followed by two 60 min periods of intravenous infusion with GLP-1 at 0.4 and 1.2 pmol·kg⁻¹·min⁻¹ in the interdigestive experiments. During the initial 10 min of each infusion period, GLP-1 was infused at double doses (0.8 and 2.4 pmol·kg⁻¹·min⁻¹, respectively) to quickly establish steady state plasma levels. On the postprandial study day, the same intravenous infusions were administered with an intraduodenal

25

30

lipid perfusion at 2.5 kcal/min (2.5 ml/min, Lipofundin MCT 10%, Braun Co., Melsungen, Germany). This lipid preparation consisted of 50% medium-chain triglycerides (MCT) and 50% long-chain triglycerides (LCT, soya bean oil). In the postprandial experiments, measurements continued for another 60 min with ongoing lipid perfusion after cessation of intravenous GLP-1.

Throughout the experiment, blood samples were taken immediately before the start of intravenous infusion, as well as in 10-min intervals thereafter. Blood was collected in ice-chilled EDTA tubes containing 1000 aprotinin (kallikrein inhibitory units/ml blood) and was centrifuged immediately. The plasma was stored at -20°C until assayed.

Example 2. *Motility recordings of the antro-pyloro-duodenal region*

Perfusion manometry was recorded using a nine-lumen duodenal sleeve/side-hole catheter (Dentsleeve, South Australia, Australia). The manometric assembly incorporated a 4.5 cm long sleeve sensor, two antral side-holes (2 cm apart), and three duodenal side-holes (2 cm apart), beginning at the proximal and distal end of the sleeve, respectively. Two more side-holes were positioned across the sleeve spaced 1.5 cm apart. An additional lumen located 12 cm distal to the sleeve sensor was used for duodenal perfusion.

The correct position of the duodenal probe with the sleeve array straddling the pylorus was fluoroscopically checked before start of each experiment, and monitored throughout each study by measuring the transmucosal potential difference at the distal antral and the proximal duodenal port, as previously described (Schirra *et al. J Clin Invest* 97, 92-103 (1996)). A difference of at least -15 mV indicated correct transpyloric position of the tube.

The motility channels were perfused at a rate of 0.3 ml/min using a low-compliance pneumohydraulic pump (Arndorfer Medical Specialists, Greendale, WI). Pressures were measured by external transducers. Data were simultaneously recorded on the screen of a multichannel chart system (PC Polygraph, Synectics Medical, Stockholm, Sweden) and stored in the memory of a personal computer. Data were sampled and digitized at 8 Hz followed by digital smoothing by a factor of 2.

Example 3. *Intravenous infusion of GLP-1 increases plasma GLP-1 levels*

Intravenous infusion of GLP-1 dose-dependently increased plasma GLP-1 in the experiments with both duodenal saline and lipid reaching constant plasma levels within 20 min after start of infusion (Fig. 1, Table 3). Steady state plasma levels amounted to 6.2 ± 0.6 pmol/l (low dose GLP-1) and 13.1 ± 0.9 pmol/l (high dose GLP-1) with duodenal saline, and to 7.1 ± 0.6 pmol/l (low dose GLP-1) and 14.4 ± 0.9 pmol/l (high dose GLP-1) with duodenal lipid, respectively. Duodenal lipid slightly but significantly increased plasma GLP-1 compared to duodenal saline perfusion (2.4 ± 0.3 pmol/l vs. 1.4 ± 0.2 pmol/l, $P < 0.05$). This difference remained constant with infusion of both loads of exogenous GLP-1. After cessation of GLP-1 during ongoing lipid perfusion, plasma GLP-1 immediately decreased and gradually returned to preinfusion values.

Table 1. Effect of physiological (low dose, $0.4 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and supraphysiological (high dose, $1.2 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) doses of GLP-1 on plasma glucose, and plasma immunoreactivities of GLP-1, insulin, glucagon, and pancreatic polypeptide with intraduodenal perfusion of saline or lipid (2.5 kcal/min , 2.5 ml/min).

	GLP-1 (pmol/l-60 min)	Plasma glucose (mmol/l-60 min)	Insulin (mU/l-60 min)	Glucagon (pg/ml-60 min)	Pancreatic polypeptide (pg/ml-60 min)
Intraduodenal Saline					
Saline iv.	-0.8 ± 1.0	0.3 ± 0.1	1.5 ± 1.7	22.4 ± 16.2	10.8 ± 29.7
GLP-1 iv.: low dose	$25.2 \pm 2.6^*$	$-2.3 \pm 0.3^*$	$11.3 \pm 2.4^*$	$-59.0 \pm 37.1^*$	$-90.5 \pm 32.7^*$
GLP-1 iv.: high dose	$67.4 \pm 4.0^{* \ddagger}$	$-2.5 \pm 0.4^*$	$11.9 \pm 5.1^*$	$-68.6 \pm 28.9^*$	$-128.9 \pm 33.2^*$
Intraduodenal Lipid					
Saline iv.	$7.5 \pm 1.7 \downarrow$	$-1.2 \pm 0.3 \downarrow$	$25.5 \pm 4.4 \downarrow$	$145.9 \pm 21.6 \downarrow$	$298.5 \pm 66.9 \downarrow$
GLP-1 iv.: low dose	$33.8 \pm 2.9^{* \downarrow}$	$-4.7 \pm 0.5^{* \downarrow}$	$24.7 \pm 4.8 \downarrow$	$99.7 \pm 25.1^{* \downarrow}$	$7.7 \pm 55.4^{* \downarrow}$
GLP-1 iv.: high dose	$75.2 \pm 3.9^{* \ddagger \downarrow}$	$-4.5 \pm 0.7^{* \downarrow}$	$31.4 \pm 8.4 \downarrow$	$48.0 \pm 33.8^{* \ddagger \downarrow}$	$-104.2 \pm 71.1^{* \ddagger \downarrow}$
Saline iv.: recovery	$17.1 \pm 3.1^{* \ddagger \downarrow}$	$-0.8 \pm 0.6^{* \ddagger \downarrow}$	18.3 ± 2.7	$189.0 \pm 44.7^{* \ddagger \downarrow}$	$294.4 \pm 93.0^{* \ddagger \downarrow}$

Mean \pm SEM of AUC over basal during each 60 min infusion period. N=11. For the same intraduodenal perfusion: *: $P < 0.05$ vs. saline iv., \ddagger : $P < 0.05$ vs. low dose GLP-1, \downarrow : $P < 0.05$ vs. high dose GLP-1. For different intraduodenal perfusions: \downarrow : $P < 0.05$ vs. the same intravenous infusion/intraduodenal saline.

Example 4. GLP-1 inhibits antro-duodenal motility

Transmucosal potential difference indicated correct transpyloric position of the probe during $95.9 \pm 1.6\%$ and $92.5 \pm 2.0\%$ of recording time with duodenal saline and lipid, respectively.

In the interdigestive experiments with duodenal saline perfusion, both dosages of GLP-1 significantly inhibited number, amplitudes, and motility indexes of contractions in the antrum

and also in the duodenum (Fig. 2A, Table 1). The inhibition of antro-duodenal motility occurred within 2 min after starting the low dose of GLP-1. Within 20 min, the low dose GLP-1 nearly completely inhibited antro-duodenal motility. Even the low dose GLP-1 completely abolished the antral waves propagating across the pylorus to the duodenum over a distance of 4.5 cm and 6.5 cm, respectively (Table 2).

Compared to duodenal saline, duodenal lipid perfusion inhibited antral and duodenal contractility. Against a background of duodenal lipid, exogenous GLP-1 dose-dependently diminished number, amplitude, and motility indexes of antral and duodenal contractions to a residual level comparable to the interdigestive state (Fig. 2B, Table 1). With cessation of GLP-1, antro-duodenal motility immediately increased and returned to preinfusion activity. The duration of contractions remained unchanged with GLP-1 in all experiments.

Table 2. Effect of physiological (low dose, $0.4 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and supraphysiological (high dose, $1.2 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) doses of GLP-1 on antral and duodenal motility with intraduodenal perfusion of saline or lipid (2.5 kcal/min, 2.5 ml/min).

	Antral Motility Contractions (number / 60 min)	Motility Index (mmHg*s/60 min)	Amplitude (mmHg)	Duodenal Motility Contractions (number / 60 min)	Motility Index (mmHg*s/60 min)	Amplitude (mmHg)
Intraduodenal						
Saline						
Saline iv.	69.3 ± 9.9	18601 ± 5758	63.0 ± 7.5	183.7 ± 25.1	19502 ± 4321	39.5 ± 3.6
GLP-1 iv.:						
low dose	$12.6 \pm 2.8^*$	$1478 \pm 564^*$	$31.1 \pm 3.1^*$	$36.0 \pm 14.7^*$	$3447 \pm 1624^*$	$25.0 \pm 2.5^*$
high dose	$9.6 \pm 2.8^*$	$957 \pm 322^*$	$21.1 \pm 3.1^{* \ddagger}$	$16.6 \pm 3.9^*$	$1119 \pm 226^*$	$17.5 \pm 1.9^{* \ddagger}$
Intraduodenal						
Lipid						
Saline iv.	$24.3 \pm 4.5 \text{ !} \#$	$2749 \pm 695 \text{ !}$	$32.5 \pm 3.1 \text{ !}$	$85.4 \pm 16.4 \text{ !} \#$	$9545 \pm 2891 \text{ !} \#$	$32.9 \pm 3.1 \text{ !} \#$
GLP-1 iv.:						
low dose	$12.1 \pm 4.1^*$	$880 \pm 351^*$	$23.6 \pm 3.1^*$	$29.5 \pm 5.8^*$	$2776 \pm 890^*$	$26.0 \pm 3.3^*$
high dose	$9.2 \pm 2.9^*$	$628 \pm 249^*$	$23.0 \pm 2.7^*$	$16.6 \pm 5.9^{* \ddagger}$	$1174 \pm 499^{* \ddagger}$	$19.2 \pm 3.1^{* \ddagger}$
Saline iv.:						
recovery	$28.7 \pm 7.5 \text{ \ddagger \#}$	$3147 \pm 785 \text{ \ddagger \#}$	$34.7 \pm 3.8 \text{ \ddagger \#}$	$69.6 \pm 21.2 \text{ \ddagger \#}$	$8732 \pm 3574 \text{ \ddagger \#}$	$36.6 \pm 4.5 \text{ \ddagger \#}$

Mean \pm SEM of actual values during each 60 min infusion period. Values for contractions and motility indices represent the sum of two antral and three duodenal side holes, respectively. N=1. For the same intraduodenal perfusion: *: $P < 0.05$ vs. saline iv., \ddagger : $P < 0.05$ vs. low dose GLP-1, $\#$: $P < 0.05$ vs. high dose GLP-1. For different intraduodenal perfusions: !: $P < 0.05$ vs. saline iv./intraduodenal saline, #: $P < 0.05$ vs. low dose GLP-1/intraduodenal saline.

5

Table 3. Effect of physiological (low dose, $0.4 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and supraphysiological (high dose, $1.2 \text{ pmol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) doses of GLP-1 on pyloric motility and antro-pyloro-duodenal wave propagation with intraduodenal perfusion of saline or lipid (2.5 kcal/min, 2.5 ml/min).

	Pyloric Motility IPPW (number / 10 min)	Pyloric tone (mmHg)	Antro-Pyloro-Duodenal Propagated Contractions (number / 60 min)
Intraduodenal Saline			
Saline iv.	1.6 ± 0.3	0.2 ± 0.5	9.8 ± 1.3
GLP-1 iv.: low dose	$10.1 \pm 1.9^*$	$5.3 \pm 0.9^*$	0.0^*
GLP-1 iv.: high dose	$8.6 \pm 1.3^*$	$7.3 \pm 1.5^{* \ddagger}$	0.0^*
Intraduodenal Lipid			
Saline iv.	$14.7 \pm 1.5 \text{ ; } \#$	$3.1 \pm 0.4 \text{ ; } \#$	-
GLP-1 iv.: low dose	15.7 ± 2.5	$7.4 \pm 1.0^{* \#}$	-
GLP-1 iv.: high dose	15.5 ± 2.1	$8.6 \pm 1.0^*$	-
Saline iv.: recovery	$6.2 \pm 1.1^{* \ddagger \#}$	$5.1 \pm 0.8^{* \ddagger \#}$	-

Mean \pm SEM of actual values (IPPW) and of values over basal (tone) during each 60 min infusion period. N=11. For the same intraduodenal perfusion: *: $P < 0.05$ vs. Saline iv., \ddagger : $P < 0.05$ vs. low dose GLP-1, $\#$: $P < 0.05$ vs. high dose GLP-1. For different intraduodenal perfusions: ;: $P < 0.05$ vs. saline iv./intraduodenal saline, #: $P < 0.05$ vs. low dose GLP-1/intraduodenal saline. IPPW: isolated pyloric pressure waves.

10 Example 5. *GLP-1 stimulates pyloric motility*

Pyloric tone increased in a dose-dependent fashion with exogenous GLP-1 in the interdigestive state (Fig. 3B, Table 2). Even with low dose GLP-1, the effect was significant within 10 min and fully established within 20 min after starting GLP-1 infusion. Pyloric tone stably increased with duodenal lipid when compared to duodenal saline, which was further enhanced by GLP-1 addition (Fig. 3B, Table 2). Pyloric tone steadily decreased after cessation of GLP-1, parallel to declining plasma levels. During the last 30 minutes of the experiment, pyloric tone was not significantly different from the first 60 min without GLP-1, indicating

complete recovery (3.9 ± 0.7 mmHg for the last 30 min vs. 3.1 ± 0.4 mmHg during the first 60 min of the experiments, $P=0.29$).

IPPWs immediately and significantly increased with infusion of GLP-1 (Fig. 3A, Table 2). In contrast to the tonic pyloric response, however, increases of IPPWs were not dose-dependent and exhibited a fast increase followed by a marked decline. The latter phenomenon held true for both lipid alone and GLP-1 against a background of duodenal lipid. Stimulation of IPPW with duodenal lipid was significantly stronger than with GLP-1 alone, and duodenal lipid and exogenous GLP-1 did not act in an additive manner. Cessation of GLP-1 significantly decreased IPPWs compared to measurements in the initial 60 min without GLP-1.

GLP-1 triggered quickly duodenal phase III-like episodes in most of the volunteers in the interdigestive state. Duodenal phase III-like activity has been previously demonstrated in response to a variety of stimuli such as intraduodenal infusion of dextrose (Hedde *et al.*, *Gut* 29, 1349-57 (1988)) and lipid (Hedde *et al.*, *Am J Physiol* 254, G671-9 (1988)), intravenous β -endorphin (Camilleri *et al.*, *Am J Physiol* 251, G147-54 (1986)), during stress induced by cold (Fone *et al.*, *Gastroenterol.* 98, 1155-61 (1990)), pain (Thompson *et al.*, *Gastroenterol.* 83, 1200-6 (1982)), or labyrinthine stimulation (Stanghellini *et al.*, *Gastroenterol.* 85, 83-91 (1983)), and acute hyperglycaemia (Fraser ET AL., *Gut* 32, 475-8 (1991)). Interestingly, intraduodenal lipid stimulated duodenal phase III-like episodes in the present study. However, GLP-1 infusion in conjunction with lipid perfusion was accompanied by duodenal phases III in only two out of 11 subjects. All duodenal phase III-like episodes strongly paralleled the initiation of pyloric stimulation, either by GLP-1 or by duodenal lipid. Therefore, the initial phasic and transient stimulation of the duodenum may originate in the pylorus, and, although its functional significance is still uncertain, it may indicate a rather unspecific reaction in response to a strong stimulation of the pyloric muscle by GLP-1 or lipid. Initiation of phasic and tonic stimulation of the pylorus was almost paralleled by duodenal phases III with both doses of GLP-1 in the interdigestive state, occurring within 10 min after GLP-1 infusion (fig. 4). These duodenal phases III were demonstrated in seven out of 11 volunteers after the low dose of GLP-1, and in five volunteers after the high dose of GLP-1. Seven out of 11 volunteers immediately developed a duodenal phase III in parallel with pyloric stimulation after intraduodenal lipid infusion. The typical pattern of the antro-pyloro-duodenal motility seen with GLP-1 infusion is shown in Figure 5.

Example 6. GLP-1 decreases plasma glucose

In the interdigestive experiments, basal plasma glucose levels were 4.3 ± 0.1 mmol/l and significantly dropped with both doses of GLP-1 (fig. 5A, Table 3). Plasma glucose levels decreased slightly compared to duodenal saline perfusion with duodenal lipid. This decrease was further pronounced with both doses of GLP-1, followed by a significant increase after cessation of GLP-1. However, nadir of plasma glucose amounted to 3.58 ± 0.12 mmol/l and 3.64 ± 0.14 mmol/l at 20 min each with low and high doses of GLP-1, respectively. These nadir values do not represent hypoglycemia.

Basal plasma levels of insulin averaged 6.0 ± 0.7 mU/l and 6.1 ± 0.7 mU/l before start of duodenal saline and lipid infusion, respectively. Both doses of GLP-1 led to a short, initial rise of insulin, followed by a decline (Fig. 5C). Compared to duodenal saline, duodenal lipid elicited a small, but significant increase in plasma insulin, which was maintained throughout each study (Table 3).

Basal levels of glucagon amounted to 91.5 ± 8.8 pg/ml and 88.2 ± 8.6 pg/ml in the studies with duodenal saline and lipid, respectively (difference not significant). With duodenal saline and lipid, GLP-1 significantly and dose-dependently diminished plasma glucagon (Fig. 5B). Plasma glucagon then immediately increased with cessation of GLP-1 in the lipid experiments. Compared to saline perfusion, duodenal lipid significantly raised glucagon, which then remained elevated throughout the experiments (Table 3).

Example 7. GLP-1 dose-dependently decreases pancreatic polypeptide levels

The inhibition of pancreatic polypeptide (PP), a hormone of the endocrine pancreas, is under strong vagal cholinergic control. PP is inhibited in humans with subcutaneous and intravenous GLP-1 after oral meal ingestion (Schirra *et al. J. Endocrinol.* 156, 177-86 (1998); Schirra *et al. Proc Assoc Am Physicians* 109, 84-97 (1997); Dupre *et al. Diabetes* 44, 626-30 (1995)). Intestinal stimulation of PP release requires stimulation of enteropancreatic cholinergic reflexes by duodenal delivery of nutrients (Schwartz, *Gastroenterol.* 85, 1411-25 (1983)), and a reduced duodenal nutrient load during retarded gastric emptying would explain this PP response. In the results marshaled by the inventor, PP release triggered by lipid perfusion directly into the duodenum dose-dependently decreased with GLP-1 infusion, followed by a prompt and complete

recovery after cessation of GLP-1 treatment. Moreover, in the interdigestive studies, PP was significantly reduced below basal levels. Therefore, the present inventor finds that GLP-1 inhibits efferent vagal-cholinergic activity, thereby diminishing PP release and at least partially contributing to the inhibition of antral and duodenal motility via a central pathway. Receptors for GLP-1 are present in circumventricular organs like the subfornical organ, the nucleus of the solitary tract and the area postrema (Göke *et al.*, *Eur. J. Neurosci* 7, 2294-2300 (1995)). In addition, it has been recently shown in rat that GLP-1 induced inhibition of gastric emptying involves a capsaicin-sensitive pathway indicating an interaction with vagal afferent nerves (Imeryuz *et al.*, *Am. J. Physiol.* 273, G920-7 (1997)). A direct action of GLP-1 on pancreatic PP cells or a paracrine effect via somatostatin seems unlikely because GLP-1 induces stimulation, instead of inhibition of PP release from isolated human pancreatic islets (Fehmann *et al.*, *Pancreas* 11, 196-200 (1995)).

Basal levels of pancreatic polypeptide averaged 64.3 ± 8.6 pg/ml and 71.9 ± 8.7 pg/ml before start of duodenal saline and lipid, respectively ($P=0.32$). With duodenal lipid, levels of pancreatic polypeptide markedly rose when compared to saline perfusion (Fig. 5D, table 3). Intravenous GLP-1 dose-dependently diminished pancreatic polypeptide with and without duodenal lipid, and pancreatic polypeptide was significantly reduced below basal levels in the experiments with duodenal saline. After cessation of GLP-1, an instantaneous increase of pancreatic polypeptide levels was observed in the studies with duodenal lipid perfusion, indicating a complete recovery.

CLAIMS

1. A method for inhibiting antro- duodenal motility in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a GLP-1 molecule.
5
2. The method for inhibiting antro- duodenal motility in claim 1, wherein said GLP-1 molecule is a GLP-1(7-36)amide.
3. A dosage unit for use in inhibiting antro- duodenal motility in a patient comprising GLP-
10 1 and a pharmaceutically suitable excipient.
4. The dosage unit of claim 3, wherein dose of said GLP-1 molecule is in the range of about 0.4-2.4 pmol·kg⁻¹·min⁻¹.
- 15 5. The dosage unit of claim 4, wherein dose of said GLP-1 molecule is in the range of about 0.8-1.2 pmol·kg⁻¹·min⁻¹.
6. The dosage unit of claim 3, wherein said pharmaceutically suitable excipient is orally, parenterally or subcutaneously acceptable.
20
7. The dosage unit of claim 3, wherein said pharmaceutically suitable excipient is intravenously acceptable.
8. A method for premedicating in endoscopic procedures, which comprises administering to
25 the patient a therapeutically effective amount of a GLP-1 molecule.
9. The method for premedicating in endoscopic procedures in claim 8, wherein said GLP-1 molecule is a GLP-1(7-36)amide.

10. A method for treating or preventing non-infectious acute or chronic diarrhea in a patient in need thereof which comprises administering to the patient a therapeutically effective amount of a GLP-1 molecule.
- 5 11. The method for treating or preventing non-infectious acute and chronic diarrhea in claim 10, wherein said GLP-1 molecule is a GLP-1(7-36)amide.
12. A method for treating or preventing postoperative dumping syndrome in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a
10 GLP-1 molecule.
- 13 The method for treating or preventing postoperative dumping syndrome in claim 12, wherein said GLP-1 molecule is a GLP-1(7-36)amide.
- 15 14. A method for treating or preventing irritable bowel syndrome in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a GLP-1 molecule.
15. The method for treating or preventing irritable bowel syndrome in claim 14, wherein said
20 GLP-1 molecule is a GLP-1(7-36)amide.
16. A method for treating or preventing symptoms associated with narcotics withdrawal in a patient in need thereof, which comprises administering to the patient a therapeutically effective amount of a GLP-1 molecule.
25
17. The method for treating or preventing symptoms associated with narcotics withdrawal in claim 16, wherein said GLP-1 molecule is a GLP-1(7-36)amide.
- 30 18. A method according to claim 1, wherein said GLP-1 molecule is administered orally or parenterally.


19. Use of a GLP-1 molecule for the preparation of a medicament for inhibiting antro-duodenal motility, for premedicating in endoscopic procedures, for treating or preventing non-infectious acute or chronic diarrhea, for treating or preventing post-operative dumping syndrome, for treating or preventing irritable bowel syndrome, or for treating or preventing symptoms
5 associated with narcotics withdrawal.

20. The use in claim 19 wherein said GLP-1 molecule is a GLP-1(7-36)amide.

21. The use in claims 19 or 20, wherein said medicament is to be administered orally or
10 parenterally.


1/10

FIG. 1




SUBSTITUTE SHEET (RULE 26)

2/10

FIG. 2A

SUBSTITUTE SHEET (RULE 26)

3/10

FIG. 2B


SUBSTITUTE SHEET (RULE 26)

4/10

FIG. 3A


ISOLATED PYLORIC PRESSURE WAVES

← DUODENAL PERFUSION OF SALINE →




SUBSTITUTE SHEET (RULE 26)

5/10


FIG. 3B

SUBSTITUTE SHEET (RULE 26)

6/10
FIG. 4


SUBSTITUTE SHEET (RULE 26)

7/10





SUBSTITUTE SHEET (RULE 26)

8/10





SUBSTITUTE SHEET (RULE 26)

9/10

FIG. 6A**FIG. 6C**

SUBSTITUTE SHEET (RULE 26)

10/10

FIG. 6B**FIG. 6D**

SUBSTITUTE SHEET (RULE 26)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68112 A3

(51) International Patent Classification⁷: **A61K 38/26**,
A61P 1/12, 1/00, 41/00, 25/30

(21) International Application Number: PCT/EP01/02882

(22) International Filing Date: 14 March 2001 (14.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/189,091 14 March 2000 (14.03.2000) US

(71) Applicants and

(72) Inventors: **GÖKE, Burkhard** [DE/DE]; Römerstrasse
47 a, 82131 Gauting (DE). **SCHIRRA, Jörg** [DE/DE];
Meyerbeerstrasse 4, 81247 München (DE).

(74) Agent: **DÖRRIES, FRANK-MOLNIA & POHLMAN**;
Postfach 221661, 80538 Munich (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

(88) Date of publication of the international search report:

7 March 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: EFFECTS OF GLUCAGON-LIKE PEPTIDE-1 (7-36) ON ANTRO-PYLORO-DUODENAL MOTILITY



(57) Abstract: The present invention provides an effective method for inhibiting antro-duodenal motility in healthy subjects and patients suffering from various disorders, without the side effects associated with other pharmaceutical compositions. GLP-1 slows antro-duodenal motility and may be used for the treatment or prevention of gastrointestinal disorders such as diarrhea, postoperative dumping syndrome and irritable bowel syndrome, and also premedication in endoscopic procedures.

WO 01/68112 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/02882

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/26 A61P1/12 A61P1/00 A61P41/00 A61P25/30

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, MEDLINE, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	SCHIRRA J ET AL: "Effects of glucagon-like peptide-1(7-36)amide on antro-pyloro-duodenal motility in the interdigestive state and with duodenal lipid perfusion in humans." GUT, vol. 46, no. 5, May 2000 (2000-05), pages 622-631, XP001013031 ISSN: 0017-5749 the whole document ----- -/--	1-21



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

16 October 2001

Date of mailing of the international search report

31/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Valcarcel, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/02882

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category "	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHIRRA J ET AL: "Effects of GLP-1 on pyloric motility and endocrine pancreatic response in the interdigestive state and with duodenal lipid." GASTROENTEROLOGY, vol. 116, no. 4 PART 2, April 1999 (1999-04), page A643 XP001008660 Digestive Disease Week and the 100th Annual Meeting of the American Gastroenterological Association; Orlando, Florida, USA; May 16-19, 1999 ISSN: 0016-5085	1-7, 18-21
Y	the whole document ----	8-17
X	SCHIRRA J ET AL: "Endogenous GLP-1(7-36)amide controls endocrine pancreatic secretion and antroduodenal motility in human." GASTROENTEROLOGY, vol. 114, no. 4 PART 2, 15 April 1998 (1998-04-15), page A1178 XP001013003 Digestive Diseases Week and the 99th Annual Meeting of the American Gastroenterological Association; New Orleans, Louisiana, USA; May 16-22, 1998 ISSN: 0016-5085	1-7, 18-21
Y	the whole document ----	8-17
X	SCHIRRA J ET AL: "Differential effects of subcutaneous GLP - 1 on gastric emptying, antroduodenal motility, and pancreatic function in men." PROCEEDINGS OF THE ASSOCIATION OF AMERICAN PHYSICIANS, (1997 JAN) 109 (1) 84-97., XP001013311	1-7, 18-21
Y	the whole document ----	8-17
X	SCHIRRA J ET AL: "Effects of GLP-1 on human antro-pyloro-duodenal motility." GASTROENTEROLOGY, vol. 110, no. 4 SUPPL., 1996, page A1116 XP001012953 96th Annual Meeting of the American Gastroenterological Association and the Digestive Disease Week; San Francisco, California, USA; May 19-22, 1996 ISSN: 0016-5085	1-7, 18-21
Y	the whole document ----- -/--	8-17

INTERNATIONAL SEARCH REPORT

Inter: International Application No

PCT/EP 01/02882

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHIRRA J ET AL: "Differential effects of subcutaneous GLP-1 on gastric emptying, insulin release and exocrine pancreatic secretion in man." GASTROENTEROLOGY, vol. 108, no. 4 SUPPL., 1995, page A1003 XP001013040 95th Annual Meeting of the American Gastroenterological Association and Digestive Disease Week; San Diego, California, USA; May 14-17, 1995 ISSN: 0016-5085	1-7, 18-21
Y	the whole document	8-17
Y	TOLESSA TESFAYE ET AL: "Inhibitory effect of glucagon-like peptide-1 on small bowel motility. Fasting but not fed motility inhibited via nitric oxide independently of insulin and somatostatin." JOURNAL OF CLINICAL INVESTIGATION, vol. 102, no. 4, 15 August 1998 (1998-08-15), pages 764-774, XP001012974 ISSN: 0021-9738 the whole document	1-21
Y	GIRALT M ET AL: "Sympathetic pathways mediate GLP-1 actions in the gastrointestinal tract of the rat." REGULATORY PEPTIDES, vol. 74, no. 1, 24 April 1998 (1998-04-24), pages 19-25, XP001013010 ISSN: 0167-0115 the whole document	1-21
Y	MOSE P L ET AL: "Suppression of duodenal motility during ERCP by L-hyoscyamine versus glucagon: A randomized, prospective, double blinded trial." GASTROENTEROLOGY, vol. 114, no. 4 PART 2, 15 April 1998 (1998-04-15), page A533 XP001013004 Digestive Disease Week and the 99th Annual Meeting of the American Gastroenterological Association; New Orleans, Louisiana, USA; May 16-22, 1998 ISSN: 0016-5085 the whole document	8,9

	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/02882

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NASLUND ERIK ET AL: "Distal small bowel hormones: Correlation with fasting antroduodenal motility and gastric emptying." DIGESTIVE DISEASES AND SCIENCES, vol. 43, no. 5, May 1998 (1998-05), pages 945-952, XP001013043 ISSN: 0163-2116	1
Y	the whole document	2-21
Y	ANVARI MEHRAN ET AL: "Effects of GLP-1 on gastric emptying, antropyloric motility, and transpyloric flow in response to a nonnutrient liquid." DIGESTIVE DISEASES AND SCIENCES, vol. 43, no. 6, June 1998 (1998-06), pages 1133-1140, XP001013009 ISSN: 0163-2116 the whole document	1-21
X	ANVARI M ET AL: "Sites of action of GLP-1 to inhibit transpyloric flow and gastric emptying of non-nutrient liquids in conscious dogs." CLINICAL AND INVESTIGATIVE MEDICINE, vol. 18, no. 4 SUPPL., 1995, page 856 XP001013012 Annual Meeting of the Canadian Society for Clinical Investigation and the Royal College of Physicians and Surgeons of Canada;Montreal, Quebec, Canada; September 13-17, 1995 ISSN: 0147-958X	12,13,19
Y	the whole document	1-11, 14-18, 20,21
Y	ANVARI M ET AL: "Effect of GLP-1 on antropyloric motility, transpyloric flow and gastric emptying of non-nutrient liquids in conscious dogs." GASTROENTEROLOGY, vol. 108, no. 4 SUPPL., 1995, page A563 XP001012952 95th Annual Meeting of the American Gastroenterological Association and Digestive Disease Week;San Diego, California, USA; May 14-17, 1995 ISSN: 0016-5085 the whole document	1-21
X	WO 99 64060 A (EFENDIC SUAD ;HELLSTROEM PER (SE)) 16 December 1999 (1999-12-16)	1-7,14, 15,18-21
Y	the whole document	8-13,16, 17

	--- -/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/02882

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 39022 A (1149336 ONTARIO INC) 11 September 1998 (1998-09-11) The whole document (see specially claim 1) ----	16,17
A	VECHT J ET AL: "The dumping syndrome. Current insights into pathophysiology, diagnosis and treatment." SCANDINAVIAN JOURNAL OF GASTROENTEROLOGY. SUPPLEMENT, (1997) 223 21-7. REF: 95, XP001013316 cited in the application the whole document ----	12,13
Y	QUIGLEY E.M.M.: "Gastroduodenal motility." CURRENT OPINION IN GASTROENTEROLOGY, (1999) 15/6 (481-491)., , XP001013511 page 484, left-hand column, paragraph 3 ----	1-7, 18-21
Y	NAUCK M A: "Is glucagon - like peptide 1 an incretin hormone?." DIABETOLOGIA, (1999 MAR) 42 (3) 373-9. REF: 67, XP001013315 the whole document ----	1-21
Y	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; Scandinavian Journal of Gastroenterology, 1979 QVIGSTAD T ET AL: "Comparison of Glucagon Atropine and Placebo as Pre Medication for Endoscopy of the Upper Gastro Intestinal Tract" Database accession no. PREV197968075464 XP002174496 abstract ----	8,9
Y	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; Scandinavian Journal of Gastroenterology, 1995 CHANG F-Y ET AL: "A randomized study comparing glucagon and hyoscine N-butyl bromide before endoscopic retrograde cholangiopancreatography" Database accession no. PREV199598218113 XP002174497 abstract -----	8,9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/02882

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9964060	A	16-12-1999	AU 4669799 A	30-12-1999
			EP 1094834 A1	02-05-2001
			NO 20006194 A	25-01-2001
			PL 343104 A1	30-07-2001
			WO 9964060 A1	16-12-1999
WO 9839022	A	11-09-1998	US 5846937 A	08-12-1998
			AU 6286698 A	22-09-1998
			WO 9839022 A1	11-09-1998

